

Problem Set 7: Due Monday, September 29

Problem 1: In each of the following cases, determine if the given function $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation. If yes, explain why. If no, provide an explicit counterexample.

- a. $T\left(\begin{pmatrix} x \\ y \end{pmatrix}\right) = \begin{pmatrix} xy \\ x + y \end{pmatrix}$
b. $T\left(\begin{pmatrix} x \\ y \end{pmatrix}\right) = \begin{pmatrix} y \sin^2(x^3) + y \cos^2(x^3) \\ y \end{pmatrix}$
c. $T\left(\begin{pmatrix} x \\ y \end{pmatrix}\right) = \begin{pmatrix} 2x + 3y \\ 1 + y \end{pmatrix}$

Problem 2: Consider the linear transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by

$$T\left(\begin{pmatrix} x \\ y \end{pmatrix}\right) = \begin{pmatrix} -y \\ -x \end{pmatrix}.$$

Plot the values of at least 4 points and where T sends them, and then use that to describe the action of T geometrically.

Problem 3: Show that the linear transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by

$$T\left(\begin{pmatrix} x \\ y \end{pmatrix}\right) = \begin{pmatrix} x + 2y \\ 3x + 6y \end{pmatrix}$$

is not injective.

Problem 4: Consider the linear transformation $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by

$$T\left(\begin{pmatrix} x \\ y \end{pmatrix}\right) = \begin{pmatrix} 2x - y \\ -5x + 3y \end{pmatrix}.$$

Show that

$$\begin{pmatrix} -18 \\ 47 \end{pmatrix} \in \text{range}(T)$$

by explicitly finding $\vec{v} \in \mathbb{R}^2$ with

$$T(\vec{v}) = \begin{pmatrix} -18 \\ 47 \end{pmatrix}.$$

Problem 5: Suppose that $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $S: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ are both linear transformations. Show that $T \circ S: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is a linear transformation.