

## Writing Assignment 4: Due Wednesday, February 27

**Problem 1:** Let  $\vec{u}, \vec{w} \in \mathbb{R}^2$ . Suppose that  $\vec{w} \in \text{Span}(\vec{u})$  and  $\vec{w} \neq \vec{0}$ . Show that  $\text{Span}(\vec{w}) = \text{Span}(\vec{u})$ .

**Problem 2:** Let  $\vec{u}_1, \vec{u}_2, \vec{u}_3 \in \mathbb{R}^2$ . Show that at least one of the  $\vec{u}_i$  is in the span of the other two. That is show that either  $\vec{u}_1 \in \text{Span}(\vec{u}_2, \vec{u}_3)$ , or  $\vec{u}_2 \in \text{Span}(\vec{u}_1, \vec{u}_3)$ , or  $\vec{u}_3 \in \text{Span}(\vec{u}_1, \vec{u}_2)$ .

**Problem 3:** Let  $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be a linear transformation. Recall that

$$\text{range}(T) = \{\vec{w} \in \mathbb{R}^2 : \text{There exists } \vec{v} \in \mathbb{R}^2 \text{ with } \vec{w} = T(\vec{v})\}.$$

Notice that  $\vec{0} \in \text{range}(T)$  because we know that  $T(\vec{0}) = \vec{0}$  by Proposition 2.4.2.

- a. Show that if  $\vec{w}_1, \vec{w}_2 \in \text{range}(T)$ , then  $\vec{w}_1 + \vec{w}_2 \in \text{range}(T)$ .
- b. Show that if  $\vec{w} \in \text{range}(T)$  and  $c \in \mathbb{R}$ , then  $c\vec{w} \in \text{range}(T)$ .